

University of Groningen

Nonbullous pemphigoid

Lamberts, Aniek; Meijer, Joost M; Jonkman, Marcel F

Published in:
Journal of the American Academy of Dermatology

DOI:
[10.1016/j.jaad.2017.10.035](https://doi.org/10.1016/j.jaad.2017.10.035)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Final author's version (accepted by publisher, after peer review)

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Lamberts, A., Meijer, J. M., & Jonkman, M. F. (2018). Nonbullous pemphigoid: A systematic review. *Journal of the American Academy of Dermatology*, 78(5), 989-+. <https://doi.org/10.1016/j.jaad.2017.10.035>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Accepted Manuscript

Nonbullous cutaneous pemphigoid: a systematic review

Aniek Lamberts, MD, Joost M. Meijer, MD, Marcel F. Jonkman, MD, PhD

PII: S0190-9622(17)32593-8

DOI: [10.1016/j.jaad.2017.10.035](https://doi.org/10.1016/j.jaad.2017.10.035)

Reference: YMJD 12085

To appear in: *Journal of American Dermatology*

Received Date: 8 July 2017

Revised Date: 17 October 2017

Accepted Date: 20 October 2017

Please cite this article as: Lamberts A, Meijer JM, Jonkman MF, Nonbullous cutaneous pemphigoid: a systematic review, *Journal of American Dermatology* (2017), doi: 10.1016/j.jaad.2017.10.035.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



1 **Article type:** review

2

3 **Title:** Nonbullous cutaneous pemphigoid: a systematic review

4 Aniek Lamberts, MD¹, Joost M. Meijer, MD¹, Marcel F. Jonkman, MD, PhD¹

5

6 ¹ University of Groningen, University Medical Center Groningen, Department of
7 Dermatology, Center for Blistering Diseases, Groningen, the Netherlands

8

9 **Correspondence address**

10 Marcel F. Jonkman, MD, Ph.D.

11 Department of Dermatology

12 University Medical Center Groningen

13 Hanzeplein 1

14 9700 RB Groningen

15 The Netherlands

16

17 tel: +31-50-3612520

18 fax: +31-503612624

19 email: m.f.jonkman@umcg.nl

20

21 **Funding sources:** none

22 **IRB approval:** not applicable

23 **Conflict of interest:** MJ received a grant from Castle Creek and honoraria from
24 Roche/Genentech

25

26 **Manuscript word count:** 2292

27 **Abstract word count:** 196

28 **Conflict of interest :** MJ: shares Philae Pharmaceuticals, consultant

29 Roche/Genentech, grant Castle Creek Pharma.

30

31 **References:** 79

32 **Figures:** 1

33 **Tables:** 4

34 **Supplements:** 2

35

36 Key words: pemphigoid - nonbullous cutaneous pemphigoid – autoimmune bullous
37 disease – autoimmune blistering disease - cutaneous pemphigoid – bullous
38 pemphigoid – clinical presentation – characteristics – systematic review -
39 terminology

40 ABSTRACT

41 Background: Cutaneous pemphigoid (bullous pemphigoid) is an autoimmune bullous
42 disease that typically presents with tense bullae and severe pruritus. However,
43 bullae may be lacking, a subtype termed nonbullous cutaneous pemphigoid.

44 Objective: To summarize the reported characteristics of nonbullous cutaneous
45 pemphigoid.

46 Methods: The EMBASE and MEDLINE databases were searched using 'nonbullous
47 cutaneous pemphigoid' and various synonyms. Case reports and series describing
48 nonbullous cutaneous pemphigoid were included.

49 Results: The search identified 133 articles. After selection 39 articles were included,
50 presenting 132 cases. Erythematous, urticarial plaques (52.3%) and
51 papules/nodules (20.5%) were the most reported clinical features. The mean age at
52 presentation was 74.9 years. Histopathology was commonly nonspecific. Linear
53 depositions of IgG/C3 along the basement membrane zone were found by direct
54 immunofluorescence microscopy in 93.2%. Indirect immunofluorescence on salt split
55 skin was positive in 90.2%. The mean diagnostic delay was 22.6 months. The
56 minority of patients (9.8%) developed bullae during the reported follow-up.

57 Limitations: Results are mainly based on case reports/small case series.

58 Conclusion: Nonbullous cutaneous pemphigoid is an underdiagnosed variant of
59 pemphigoid that most often does not evolve to bullous lesions, and mimics other
60 pruritic skin diseases. Greater awareness among physicians is needed to avoid
61 delay in diagnosis.

62

63 INTRODUCTION

64 Cutaneous pemphigoid, also known as bullous pemphigoid (BP), is the most common
 65 autoimmune bullous disease affecting the skin and mucous membranes, with autoantibodies
 66 directed against the 180 kDa BP antigen (BP180) and the 230 kDa BP antigen (BP230)
 67 located in the basement membrane zone (BMZ).¹ The disease commonly affects older
 68 patients and is associated with an increased risk of mortality, as well as a significant decline
 69 in quality of life and psychological well-being.²⁻⁶

70 The clinical phenotype of cutaneous pemphigoid is polymorphic. The typical
 71 presentation consists of tense blisters that arise on erythematous, urticarial plaques, and is
 72 accompanied by severe pruritus.^{1,3} Prior to blister formation pruritus can occur as a
 73 prodrome, with or without primary skin manifestations.⁷ In contrast to the typical bullous
 74 presentation, various atypical variants of cutaneous pemphigoid have been reported with
 75 terms such as papular pemphigoid, pemphigoid nodularis, pemphigoid vegetans,
 76 erythrodermic pemphigoid, pruritic nonbullous pemphigoid and erythema multiforme-like
 77 pemphigoid.⁸⁻¹¹ The nonbullous variant of cutaneous pemphigoid presents with pruritus and
 78 various nonbullous findings on the skin, such as erythematous patches, urticarial plaques,
 79 papules, nodules, excoriations, eczema, and erythroderma. Moreover, this variant can even
 80 present without primary skin lesions, called 'pruritus on primary, non-diseased, non-inflamed
 81 skin' according to the International Clinical Classification of Itch.^{11,12} Previous articles have
 82 emphasized the need for uniform terminology and discussed the urge for renaming *bullous*
 83 *pemphigoid*, since the use of the adjective '*bullous*' is redundant considering that
 84 '*pemphigoid*' means 'resembling pemphigus' and pemphigus is Greek for '*blister*'.^{8,13-15}
 85 Borradori and Joly proposed the term '*cutaneous pemphigoid*'.¹³ To emphasize the lack of
 86 bullae in the nonbullous disease variant, we proposed to add the adjective *nonbullous*.¹⁴
 87 Hence, the terminology used in this article.

88 Cohort studies show that at least 20% of all pemphigoid patients do not have blisters
 89 at the time of diagnosis.^{3,15} Thus, nonbullous cutaneous pemphigoid is not that uncommon or
 90 atypical as may be assumed.¹⁶ The bullous and nonbullous cutaneous pemphigoid
 91 phenotypes are immunologically indistinguishable. The diagnosis is usually based on the
 92 combination of clinical presentation, histopathological findings, direct immunofluorescence

93 (DIF) microscopy, and immunoserology.¹⁵ One of the main obstacles currently is the lack of
94 consensus on the minimal diagnostic criteria of cutaneous pemphigoid.^{8,13,14,16,17} The
95 absence of blistering in nonbullous cutaneous pemphigoid can make the recognition of this
96 disease difficult for clinicians and may result in a delay of diagnosis.^{18,19}

97 The aim of our study is to characterize and define nonbullous cutaneous pemphigoid
98 by systematic review, which has not been performed before. Our study lists reported clinical
99 presentations, histopathological findings, laboratory findings, and prognosis regarding
100 patients with nonbullous cutaneous pemphigoid.

101 MATERIALS AND METHODS

102 *Search strategy*

103 The literature search for this review was conducted in the EMBASE and MEDLINE
104 databases on the 4th of November 2016. Various terms and synonyms for '*nonbullous*
105 *cutaneous pemphigoid*' were used (supplement 1). There were no limitations on article type.
106 After the selection procedure the references of all included articles were checked for missing
107 articles.

108 *Selection of articles*

109 Language was limited to Dutch, German or English. Independent screening of the titles and
110 abstracts was carried out by AL and JM. Discrepancies between the researchers were
111 resolved through discussion. All articles reporting on one or multiple cases of nonbullous
112 cutaneous pemphigoid were included. Nonbullous cutaneous pemphigoid was defined as all
113 symptomatic cases with a nonbullous phenotype, that lacked a previous history of bullae,
114 and fulfill the following diagnostic criteria of cutaneous pemphigoid: a positive DIF with linear
115 IgG and/or C3c along the BMZ and/or positive indirect immunofluorescence (IIF), in
116 combination with compatible clinical presentation, histopathological findings, or other
117 immunoserological tests. If the full text was not available online it was ordered at the national
118 library. Poster abstracts were only included if sufficient individual patient data was presented.

119 *Data collection*

120 The following variables were gathered: age at diagnosis, gender, duration of symptoms
121 before the diagnosis was made, clinical presentation, results of diagnostic tests,
122 histopathological findings, total follow-up time and whether blisters developed during follow-
123 up. Statistical analyses were done in IBM SPSS statistics 23.

124 RESULTS

125 *Systematic search results*

126 A total of 39 articles presenting a total of 132 cases of nonbullous cutaneous pemphigoid
127 were identified (supplement 2). Figure 1 displays the selection procedure. The first case of
128 nonbullous cutaneous pemphigoid was reported in 1983 by Barker et al.²⁰ The largest case
129 series was from Lamb et al.²¹, who described the clinical presentation of 53 patients
130 diagnosed with 'prodromal bullous pemphigoid'. This large case series did not present
131 individual patient characteristics concerning age, gender, duration of symptoms,
132 histopathological findings and total duration of follow-up. However, we were able to include
133 the reported clinical presentation and the number of cases that developed blisters during
134 follow-up.

135 *Clinical presentation*

136 Table 1 shows the demographics of the reported patients with nonbullous cutaneous
137 pemphigoid. The mean age at presentation was 74.9 years. The reported efflorescences and
138 configurations of skin lesions seen at dermatological examination are displayed in table 2.
139 Table 3 presents the location of skin lesions, reported in 64 of the 132 cases.

140 *Histopathology*

141 The histopathological findings were described in 53 individual cases. A perivascular infiltrate
142 was seen most frequently (n=32; 60.4%), which is a non-specific finding. Additionally, non-
143 specific findings not further specified were reported in 14 cases (26.4%). Eosinophils were
144 present in the biopsies of 25 cases (47.2%) and neutrophils in 7 cases (13.2%). Spongiosis
145 without eosinophils was reported in 10 cases (18.9%), while eosinophilic spongiosis was
146 seen in 4 patients (7.5%). The presence of dermal edema was reported in 8 cases (15.1%).
147 The presence of a microscopic subepidermal split was reported in 8 patients (15.1%).

148 *Laboratory findings*

149 Table 4 shows the reported laboratory findings of patients with nonbullous cutaneous
150 pemphigoid. In all cases DIF microscopy was performed. In cases with a negative DIF result,
151 the diagnosis was based on positive IIF with additional serological tests that specified the

152 targeted antigen. IIF was the most commonly performed immunoserological test (55 cases).
153 The substrate used in IIF was not specified in 15 cases. In the other cases monkey
154 esophagus (n=27) or human skin (n=13) were used as substrate. The BP230 ELISA was the
155 least performed immunoserological test (n=19). Additionally in four cases
156 immunoprecipitation was used to identify antigens, resulting in a positive reaction to both
157 BP180 and BP230 in one case and only a positive reaction to BP230 in three cases.
158 Eosinophilia in peripheral blood was reported in 13 of 15 cases (86.7%).

DISCUSSION

This systematic review summarizes the reported characteristics of nonbullous cutaneous pemphigoid. The most frequently reported skin efflorescences were erythematous, urticarial plaques (52.3%). Pruritus was reported in 100% of the cases. Overall, the duration between the start of symptoms and the correct diagnosis was very long (mean 22.6 months). Only 13 patients (9.8%) developed bullae during the reported follow-up, thus were actually prodromal to the bullous phase of cutaneous pemphigoid. However, in the majority of the cases (90.2%) bullae never occurred. The findings of this review show that although the clinical presentation of nonbullous cutaneous pemphigoid is various, pruritus at high age may be a clinical clue.

Our study identified several similarities in clinical characteristics of nonbullous and bullous cutaneous pemphigoid. Both present at older age (mean 74.9 versus 77.2 – 82.6 years).⁴⁻⁶ Furthermore, in both variants lesions are most frequently located on the trunk and extremities.^{18,22} Most of the skin efflorescences reported in nonbullous cutaneous pemphigoid cases can also be found in patients with cutaneous pemphigoid with bullae.^{1,15} On the other hand, mucosal involvement was rarely reported in nonbullous cutaneous pemphigoid, while reported in 10-30% of patients with cutaneous pemphigoid.^{3,15,22} In 14 cases the configurations of the skin lesions were reported to be annular, gyrate, figurate or herpetiform.^{21,23-30} Two of these patients presented with targetoid lesions.^{21,25} We also found three case reports that were possibly drug induced due to nifedipine, lisinopril and the combination of allopurinol plus colchicine.^{25,31,32} Nifedipine and lisinopril were previously associated with cutaneous pemphigoid, however it is not shown that these drugs actually cause a higher risk to develop cutaneous pemphigoid.^{33,34} Studies did show that the use of spironolactone and neuroleptics are independent risk factors for the development of cutaneous pemphigoid.^{35,36}

The reported histopathological findings in nonbullous cutaneous pemphigoid differ from cutaneous pemphigoid with typical bullae in several aspects. Histopathological findings were commonly nonspecific in nonbullous cutaneous pemphigoid and resembled eczema or prurigo nodularis. While cutaneous pemphigoid with bullae is usually characterized by the presence of eosinophilic spongiosis (>50%) and a subepidermal split (\pm 80%), in the cases

189 with nonbullous cutaneous pemphigoid histopathological findings only described eosinophilic
190 spongiosis in 7.5% and a subepidermal split in 15.1%.^{1,37} These findings emphasize the
191 need to always perform DIF microscopy and immunoserology in addition to histopathology in
192 patients in which nonbullous cutaneous pemphigoid is suspected. In nonbullous cutaneous
193 pemphigoid DIF microscopy was the most reported positive diagnostic test (positive in
194 93.2%) followed by IIF on salt-split skin (SSS) (90.2%). Both DIF microscopy and IIF on SSS
195 have a high specificity (98% and 100% respectively).³⁸ Yet, the reported percentage of
196 positive findings in DIF microscopy in nonbullous cutaneous pemphigoid might be an
197 overestimation, since this test is regarded as the reference standard for diagnosis of
198 pemphigoid and commonly the only performed immunopathological test.³⁹ Consequently the
199 diagnosis of pemphigoid might be rejected when DIF microscopy is negative and
200 immunoserological analysis might not have been performed.

201 The mean duration of symptoms of nonbullous cutaneous pemphigoid until the correct
202 diagnosis of pemphigoid was 22.6 months. These results seem to be consistent with other
203 research that also found long diagnostic delays in pemphigoid cases that lack bullae.
204 Previously, we reported a mean delay in diagnosis of 33.6 months in 15 patients with
205 nonbullous cutaneous pemphigoid.¹⁴ The studies of Zhang et al. and Sun et al. reported
206 misdiagnosis with eczema, nodular prurigo or other dermatologic diseases in all pemphigoid
207 patients that initially presented without bullae, 181 and 24 patients respectively.^{18,40} In both
208 studies the correct diagnosis was made when bullae appeared, which was after a mean
209 duration of 15.9 months and 20.75 months (range 1 month to 19 years). Although these
210 studies only identified misdiagnosis in prodromal cutaneous pemphigoid patients, they also
211 illustrate the importance of more awareness and better knowledge regarding the
212 characteristics of nonbullous cutaneous pemphigoid. In contrast, Della Torre et al. did not
213 find a significant difference in delay of diagnosis between patients with bullous (n=97) and
214 nonbullous (n=20) cutaneous pemphigoid in their cohort.³ Whether early recognition and
215 immunosuppressive treatment of nonbullous cutaneous pemphigoid can prevent later blister
216 development is unknown.

217 A much debated question is whether patients diagnosed with nonbullous cutaneous
218 pemphigoid are prodromal or have a distinct pemphigoid variant.^{10,21} The finding that the

majority of nonbullous cutaneous pemphigoid patients did not develop blisters during follow-up supports the hypothesis that nonbullous cutaneous pemphigoid is not a prodromal stage but merely a variant within the clinical spectrum of pemphigoid diseases. We can conclude that 'prodromal pemphigoid' is an incorrect term and that there is a need for consensus regarding the terminology to describe this disease variant. We strongly argue for insertion of the term nonbullous cutaneous pemphigoid in the EMTREE.

During our literature search we identified a number of other subepidermal autoimmune blistering diseases with nonbullous clinical presentations: nonbullous epidermolysis bullosa acquisita⁴¹, nonbullous linear IgA dermatosis⁴² and nonbullous pemphigoid gestationis⁴³. Furthermore we came across reports of cutaneous pemphigoid patients that first presented with bullae and later experienced a nonbullous flare-up of the disease.⁴⁴⁻⁴⁹ These cases strengthen the idea that nonbullous cutaneous pemphigoid should be seen as a disease variant within the spectrum of pemphigoid diseases. Previous publications reported a higher prevalence of BP-specific autoantibodies in older dermatology patients (>75 years) without blisters, healthy blood donors, and elderly individuals with pruritus.⁵⁰⁻⁵² How these patients fit the pemphigoid spectrum has not been clarified.

Our systematic review provides insight on reported literature on nonbullous cutaneous pemphigoid so far. A limitation of this review is that the results are mainly based on single case reports and small case series. Consequently missing values were present in the summarized data. Moreover, in some publications the clinical picture was described very briefly. A second limitation of this review is the risk of reporting bias, since cases with unusual atypical presentations are more likely to be reported in the literature. Furthermore, the finding that the majority (90.2%) of nonbullous cutaneous pemphigoid patients did not develop blisters during the reported follow-up (mean 19.8 months; range 0-72) might be slightly biased by selection, since we excluded cases of cutaneous pemphigoid that were diagnosed after bullae appeared, even though authors retrospectively described pruritic symptoms prior to blistering. However, it is uncertain whether these symptoms prior to diagnosis were caused by pemphigoid, or by other pruritic dermatoses, such as prurigo nodularis or eczema. This study therefore highlights the importance of larger observational studies with longer follow-up for a better representation of nonbullous cutaneous pemphigoid

249 Another interesting focus for future research is why patients with nonbullous
250 cutaneous pemphigoid do not develop bullae. Several factors have been suggested to
251 influence blister formation, such as autoantibody titers,⁵³ the antigens or epitopes targeted by
252 autoantibodies,^{22,54} complement involvement,^{55,56} and eosinophils.⁵⁷ More knowledge of the
253 underlying pathophysiology of this subtype of pemphigoid might lead to more awareness and
254 less delay in diagnosis of nonbullous cutaneous pemphigoid.

255 In conclusion, our review showed that the reported clinical presentation of nonbullous
256 cutaneous pemphigoid can be heterogeneous. The reported long duration of symptoms until
257 correct diagnosis (mean 22.6 months) illustrates that nonbullous cutaneous pemphigoid can
258 be difficult to recognize for clinicians. Pruritus in elderly is a common denominator in patients
259 with nonbullous cutaneous pemphigoid and in our opinion the most important clue for
260 recognition. Clinicians should therefore perform DIF on a skin biopsy and immunoserological
261 analysis on a blood sample in elderly with unexplained or refractory chronic pruritus and
262 erythematous, urticarial papules and plaques. Further study is needed to evaluate the
263 prevalence of nonbullous cutaneous pemphigoid.

274 **ABBREVIATION LIST**

275	BP	bullous pemphigoid
276	BP180	180 kDa BP antigen
277	BP230	230 kDa BP antigen
278	BMZ	basement membrane zone
279	DIF	direct immunofluorescence
280	IIF	indirect immunofluorescence
281	SSS	salt-split skin

282 REFERENCES

- 283 1. Schmidt E., Zillikens D. Pemphigoid diseases. - *Lancet*.2013 Jan
 284 26;381(9863):320-32.doi: 10.1016/S0140-6736(12)61140-4.Epub 2012 Dec 11.
- 285 2. Kouris A, Platsidaki E, Christodoulou C, et al. Quality of life, depression, anxiety
 286 and loneliness in patients with bullous pemphigoid. A case control study. - *An Bras*
 287 *Dermatol*.2016 Sep-Oct;91(5):601-603.doi: 10.1590/abd1806-4841.20164935.
- 288 3. Della Torre R, Combescure C, Cortes B, et al. Clinical presentation and
 289 diagnostic delay in bullous pemphigoid: A prospective nationwide cohort. - *Br J*
 290 *Dermatol*.2012 Nov;167(5):1111-7.doi: 10.1111/j.1365-2133.2012.11108.x.
- 291 4. Marazza G, Pham H, Scharer L, et al. Incidence of bullous pemphigoid and
 292 pemphigus in switzerland: A 2-year prospective study. - *Br J Dermatol*.2009
 293 Oct;161(4):861-8.doi: 10.1111/j.1365-2133.2009.09300.x.Epub 2009 May 8.
- 294 5. Langan S, Smeeth L, Hubbard R, Fleming K, Smith C, West J. Bullous
 295 pemphigoid and pemphigus vulgaris--incidence and mortality in the UK: Population
 296 based cohort study. - *BMJ*.2008 Jul 9;337:a180.doi: 10.1136/bmj.a180.
- 297 6. Joly P, Baricault S, Sparsa A, et al. Incidence and mortality of bullous pemphigoid
 298 in france. - *J Invest Dermatol*.2012 Aug;132(8):1998-2004.doi:
 299 10.1038/jid.2012.35.Epub 2012 Mar 15.
- 300 7. Asbrink E, Hovmark A. Clinical variations in bullous pemphigoid with respect to
 301 early symptoms. - *Acta Derm Venereol*.1981;61(5):417-21.
- 302 8. Cozzani E, Gasparini G, Burlando M, Drago F, Parodi A. Atypical presentations of
 303 bullous pemphigoid: Clinical and immunopathological aspects. *Autoimmun Rev*.
 304 2015;14(5):438-445.

- 305 9. Bernard P, Antonicelli F. Bullous pemphigoid: A review of its diagnosis,
306 associations and treatment. *Am J Clin Dermatol*. 2017.
- 307 10. Strohal R, Rappersberger K, Pehamberger H, Wolff K. Nonbullous pemphigoid:
308 Prodrome of bullous pemphigoid or a distinct pemphigoid variant? *J Am Acad*
309 *Dermatol*. 1993;29(2 II):293-299.
- 310 11. Bakker CV, Terra JB, Pas HH, Jonkman MF. Bullous pemphigoid as pruritus in
311 the elderly a common presentation. *JAMA Dermatol*. 2013;149(8):950-953.
- 312 12. Stander S, Weisshaar E, Mettang T, et al. Clinical classification of itch: A
313 position paper of the international forum for the study of itch. *Acta Derm Venereol*.
314 2007;87(4):291-294.
- 315 13. Borradori L, Joly P. Toward a practical renaming of bullous pemphigoid and all
316 its variants: Cutaneous pemphigoid. - *JAMA Dermatol*. 2014 Apr;150(4):459.doi:
317 10.1001/jamadermatol.2014.51.
- 318 14. Bakker C, Terra J, Jonkman M. Toward a practical renaming of bullous
319 pemphigoid and all its variants-reply. - *JAMA Dermatol*. 2014 Apr;150(4):459-60.doi:
320 10.1001/jamadermatol.2014.54.
- 321 15. Di Zenzo G, Della Torre R, Zambruno G, Borradori L. Bullous pemphigoid: From
322 the clinic to the bench. - *Clin Dermatol*. 2012 Jan-Feb;30(1):3-16.doi:
323 10.1016/j.clindermatol.2011.03.005.
- 324 16. Lipsker D, Borradori L. 'Bullous' pemphigoid: What are you? urgent need of
325 definitions and diagnostic criteria. *Dermatology*. 2010;221(2):131-134.
- 326 17. Feliciani C, Joly P, Jonkman M, et al. Management of bullous pemphigoid: The
327 european dermatology forum consensus in collaboration with the european

- academy of dermatology and venereology. - *Br J Dermatol.* 2015 Apr;172(4):867-77. doi: 10.1111/bjd.13717.
18. Zhang Y, Luo Y, Han Y, Tian R, Li W, Yao X. Non-bullous lesions as the first manifestation of bullous pemphigoid: A retrospective analysis of 181 cases. *J Dermatol.* 2017.
19. Hertl M, Schmidt T. Underrecognition of the heterogeneous clinical spectrum of bullous pemphigoid. *JAMA Dermatol.* 2013;149(8):954-955.
20. Barker DJ. Generalized pruritus as the presenting feature of bullous pemphigoid. *Br J Dermatol.* 1983;109(2):237-239.
21. Lamb PM, Abell E, Tharp M, Frye R, Deng J-. Prodromal bullous pemphigoid. *Int J Dermatol.* 2006;45(3):209-214.
22. Tanaka M, Hashimoto T, Dykes PJ, Nishikawa T. Clinical manifestations in 100 japanese bullous pemphigoid cases in relation to autoantigen profiles. *Clin Exp Dermatol.* 1996;21(1):23-27.
23. Altman K, Lloyd R, Longley J. Granuloma annulare-like presentation of bullous pemphigoid. *J Am Acad Dermatol.* 2015;72(5):AB119.
24. Amato DA, Silverstein J, Zitelli J. The prodrome of bullous pemphigoid. *Int J Dermatol.* 1988;27(8):560-563.
25. Axelrod S, Davis-Lorton MA. Erythema multiforme-like bullous pemphigoid drug eruption. *Ann Allergy Asthma Immunol.* 2010;105(5):A57-A58.
26. Ise Y, Suga Y, Okumura K, Negi O, Ishii N, Hashimoto T. Erythematous variety of bullous pemphigoid: Case report and literature review. *Acta Derm -Venereol.* 2016;96(3):412-413.

- 351 27. Kabuto M, Fujimoto N, Tanaka T. Two cases of annular erythema without
352 bullous lesions by autoimmune blistering diseases. *Mod Rheumatol*. 2015.
- 353 28. Park KY, Hyun MY, Yeo IK, Seo SJ, Hong CK. An eczema-like, pruritic,
354 nonbullous form of bullous pemphigoid. *J Dermatol Case Rep*. 2015;9(2):55-57.
- 355 29. Patel M, Gilbert E, Tzu J. Case report: Bullous pemphigoid associated with renal
356 cell carcinoma and invasive squamous cell carcinoma. *J Am Acad Dermatol*.
357 2012;66(4):AB44.
- 358 30. Tashiro H, Arai H, Hashimoto T, Takezaki S, Kawana S. Pemphigoid nodularis:
359 Two case studies and analysis of autoantibodies before and after the development
360 of generalized blistering. *J Nippon Med Sch*. 2005;72(1):60-65.
- 361 31. Liu VT, Cheng KY, Yu P. Where are the blisters? A case of bullous pemphigoid
362 without bullae. *J Gen Intern Med*. 2014;29:S470.
- 363 32. Ameen M, Harman KE, Black MM. Pemphigoid nodularis associated with
364 nifedipine [11]. *Br J Dermatol*. 2000;142(3):575-577.
- 365 33. Shachar E, Bialy-Golan A, Srebrnik A, Brenner S. "Two-step" drug-induced
366 bullous pemphigoid. *Int J Dermatol*. 1998;37(12):938-939.
- 367 34. Kalinska-Bienias A, Rogozinski TT, Wozniak K, Kowalewski C. Can pemphigoid
368 be provoked by lisinopril? *Br J Dermatol*. 2006;155(4):854-855.
- 369 35. Bastuji-Garin S, Joly P, Lemordant P, et al. Risk factors for bullous pemphigoid
370 in the elderly: A prospective case-control study. *J Invest Dermatol*. 2011;131(3):637-
371 643.
- 372 36. Bastuji-Garin S, Joly P, Picard-Dahan C, et al. Drugs associated with bullous
373 pemphigoid. A case-control study. *Arch Dermatol*. 1996;132(3):272-276.

- 374 37. Chan YC, Sun YJ, Ng PP, Tan SH. Comparison of immunofluorescence
375 microscopy, immunoblotting and enzyme-linked immunosorbent assay methods in
376 the laboratory diagnosis of bullous pemphigoid. *Clin Exp Dermatol.* 2003;28(6):651-
377 656.
- 378 38. Sardy M, Kostaki D, Varga R, Peris K, Ruzicka T. Comparative study of direct
379 and indirect immunofluorescence and of bullous pemphigoid 180 and 230 enzyme-
380 linked immunosorbent assays for diagnosis of bullous pemphigoid. *J Am Acad*
381 *Dermatol.* 2013;69(5):748-753.
- 382 39. Bagci IS, Horvath ON, Ruzicka T, Sardy M. Bullous pemphigoid. *Autoimmun*
383 *Rev.* 2017;16(5):445-455.
- 384 40. Sun C, Chang B, Gu H. Non-bullous lesions as the first manifestation of bullous
385 pemphigoid: A retrospective analysis of 24 cases. *J Dermatolog Treat.*
386 2009;20(4):233-237.
- 387 41. Furukita K, Ansai S, Hida Y, Kubo Y, Arase S, Hashimoto T. A case of
388 epidermolysis bullosa acquisita with unusual clinical features. *Clin Exp Dermatol.*
389 2009;34(8):e702; e704.
- 390 42. Chaudhari S, Mobini N. Linear IgA bullous dermatosis: A rare clinicopathologic
391 entity with an unusual presentation. *J Clin Aesthetic Dermatol.* 2015;8(10):43-46.
- 392 43. Ogilvie P, Trautmann A, Dummer W, Rose C, Bröcker EB, Zillikens D.
393 Pemphigoid gestationis without blisters. *Hautarzt.* 2000;51(1):25-30.
- 394 44. Borradori L, Prost C, Wolkenstein P, Bernard P, Baccard M, Morel P. Localized
395 pretibial pemphigoid and pemphigoid nodularis. *J Am Acad Dermatol.* 1992;27(5
396 II):863-867.

- 397 45. Bourke JF, Berth-Jones J, Gawkrödger DJ, Burns DA. Pemphigoid nodularis: A
398 report of two cases. *Clin Exp Dermatol*. 1994;19(6):496-499.
- 399 46. Gallo R, Parodi A, Rebora A. Pemphigoid nodularis [1]. *Br J Dermatol*.
400 1993;129(6):744-745.
- 401 47. Mochizuki M, Fujine E, Tawada C, Kanoh H, Seishima M. Pemphigoid nodularis
402 possibly induced by etanercept. *J Dermatol*. 2013;40(7):578-579.
- 403 48. Yung CW, Soltani K, Lorincz AL. Pemphigoid nodularis. *J Am Acad Dermatol*.
404 1981;5(1):54-60.
- 405 49. Ross JS, McKee PH, Smith NP, et al. Unusual variants of pemphigoid: From
406 pruritus to pemphigoid nodularis. *J CUTANEOUS PATHOL*. 1992;19(3):212-216.
- 407 50. Meijer JM, Lamberts A, Pas HH, Jonkman MF. Significantly higher prevalence of
408 circulating bullous pemphigoid-specific IgG autoantibodies in elderly patients with a
409 nonbullous skin disorder. *Br J Dermatol*. 2015;173(5):1274-1276.
- 410 51. van Beek N, Dohse A, Riechert F, et al. Serum autoantibodies against the
411 dermal-epidermal junction in patients with chronic pruritic disorders, elderly
412 individuals and blood donors prospectively recruited. *Br J Dermatol*.
413 2014;170(4):943-947.
- 414 52. Schmidt T, Sitaru C, Amber K, Hertl M. BP180- and BP230-specific IgG
415 autoantibodies in pruritic disorders of the elderly: A preclinical stage of bullous
416 pemphigoid? *Br J Dermatol*. 2014;171(2):212-219.
- 417 53. Kawahara Y, Matsumura K, Hashimoto T, Nishikawa T. Immunoblot analysis of
418 autoantigens in localized pemphigoid and pemphigoid nodularis. *Acta Derm*
419 *Venerol*. 1997;77(3):187-190.

- 420 54. Thoma-Uszynski S, Uter W, Schwietzke S, et al. BP230- and BP180-specific
421 auto-antibodies in bullous pemphigoid. *J Invest Dermatol.* 2004;122(6):1413-1422.
- 422 55. Romeijn TR, Jonkman MF, Knoppers C, Pas HH, Diercks GF. Complement in
423 bullous pemphigoid: Results from a large observational study. *Br J Dermatol.*
424 2017;176(2):517-519.
- 425 56. Nelson KC, Zhao M, Schroeder PR, et al. Role of different pathways of the
426 complement cascade in experimental bullous pemphigoid. *J Clin Invest.*
427 2006;116(11):2892-2900.
- 428 57. de Graauw E, Sitaru C, Horn M, et al. Evidence for a role of eosinophils in blister
429 formation in bullous pemphigoid. *Allergy.* 2017.
- 430 58. Bingham EA, Burrows D, Sandford JC. Prolonged pruritus and bullous
431 pemphigoid. *Clin Exp Dermatol.* 1984;9(6):564-570.
- 432 59. Borradori L, Rybojad M, Verola O, Flageul B, Puissant A, Morel P. Pemphigoid
433 nodularis. *Arch Dermatol.* 1990;126(11):1522-1523.
- 434 60. Wolf R, Ophir J, Dechner E. Nonbullous bullous pemphigoid. *Int J Dermatol.*
435 1992;31(7):498-500.
- 436 61. Wever S, Rank C, Hornschuh B, et al. Bullous pemphigoid simulation subacute
437 simple prurigo. *Hautarzt.* 1995;46(11):789-795.
- 438 62. Jeong SJ, Lee CW. Bullous pemphigoid: Persistent lesions of
439 eczematous/urticarial erythemas. *Cutis.* 1995;56(4):225-226.
- 440 63. Cliff S, Holden CA. Pemphigoid nodularis: A report of three cases and review of
441 the literature. *Br J Dermatol.* 1997;136(3):398-401.

- 442 64. Alonso-Llamazares J, Dietrich SM, Gibson LE. Bullous pemphigoid presenting
443 as exfoliative erythroderma. *J Am Acad Dermatol*. 1998;39(5 Pt 2):827-830.
- 444 65. Alonso-Llamazares J, Rogers III RS, Oursler JR, Calobrisi SD. Bullous
445 pemphigoid presenting as generalized pruritus: Observations in six patients. *Int J*
446 *Dermatol*. 1998;37(7):508-514.
- 447 66. Scrivener Y, Heid E, Grosshans E, Cribier B, Gibson LE. Erythrodermic bullous
448 pemphigoid [4] (multiple letters). *J Am Acad Dermatol*. 1999;41(4):658-659.
- 449 67. Schmidt E, Sitaru C, Schubert B, et al. Subacute prurigo variant of bullous
450 pemphigoid: Autoantibodies show the same specificity compared with classic
451 bullous pemphigoid. *J Am Acad Dermatol*. 2002;47(1):133-136.
- 452 68. Powell AM, Albert S, Gratian MJ, Bittencourt R, Bhogal BS, Black MM.
453 Pemphigoid nodularis (non-bullous): A clinicopathological study of five cases. *Br J*
454 *Dermatol*. 2002;147(2):343-349.
- 455 69. Goel A, Balchandran C, Shenoi SD, Pai B. S. Non-bullous variant of bullous
456 pemphigoid: Role of immunofluorescence in diagnosis. *Indian J Dermatol Venereol*
457 *Leprol*. 2003;69(4):294-295.
- 458 70. Mechtel D, Prugovecki V, Knopf B. Pemphigoid nodularis (non bullous) - A case
459 report. *Aktuel Dermatol*. 2003;29(7):296-299.
- 460 71. Von Felbert V, Simon D, Braathen L, Hunziker T. Pemphigoid nodularis
461 triggered by hypereosinophilic syndrome? *Hautarzt*. 2006;57(5):434-436.
- 462 72. Yesudian PD, Shah D, Giles M, Williamson D. Nonbullous variant of pemphigoid
463 in a patient with absent C4 allotype: A lesson in immunology for dermatologists. *Br J*
464 *Dermatol*. 2009;161:97.

- 465 73. Matsudate Y, Ansai S-, Hirose K, Kubo Y, Arase S. Pemphigoid nodularis: The
466 importance of ELISA for diagnosis. *Eur J Dermatol.* 2009;19(1):83-84.
- 467 74. Safa G, Darrieux L. Nonbullous pemphigoid treated with doxycycline
468 monotherapy: Report of 4 cases. *J Am Acad Dermatol.* 2011;64(6):e116; e118.
- 469 75. McCourt C, Corry A, Walsh M, McMillan C. Generalized pruritic eruption in a
470 patient with chronic kidney disease. *Dermatol Online J.* 2010;16(4).
- 471 76. Geiss Steiner J, Trüeb RM, Mühleisen B, Kerl K, French LE, Hofbauer GF.
472 Ecthyma gangrenosum-like bullous pemphigoid. *J Invest Dermatol.* 2009;129:S20.
- 473 77. Lehman JS, Kalaaji AN, Rogers 3rd. RS, Stone RA. Pemphigoid nodularis: A
474 case report. *Cutis.* 2011;88(5):224-226.
- 475 78. Balakirski G, Merk HF, Megahed M. Bullous pemphigoid: A new look at a well-
476 known disease. *Hautarzt.* 2014;65(12):1013-1016.
- 477 79. Huet F, Karam A, Lemasson G, et al. Image gallery: Erythroderma revealing a
478 nonbullous bullous pemphigoid. *Br J Dermatol.* 2016;175(5):e136-e137.

479

480 **LEGENDS TO FIGURES**

481 **Figure 1** Study selection flow diagram

482

483

484 **TABLES**

485

486 **Table 1**

487 **Table 1.** Demographics of the reported cases of nonbullous cutaneous pemphigoid

Demographic outcome measurements			Reported in no. of cases
Mean age at presentation, in years	74.9	SD 11.8; range 39-95	78
Male cases, proportion	33 (42.3%)		78
Cases experiencing pruritus, proportion	77 (100%)		77
Cases with reported mucosal lesions, proportion	1* (7.1%)		14
Mean duration of symptoms before diagnosis, in months	22.6	SD 39.1; range 0-240	50
Cases with blister development after diagnosis, proportion	13 (9.8%)		132
- Mean duration of symptoms until blisters occurred, in months	15.9	SD 8.4; range 7.5-27	5
- Mean duration from diagnosis till blisters occurred, in months	9.6	SD 8.6; range 1-21	7
Mean total follow-up, in months	19.6	SD 18.6; range 0-72	46

488 SD, standard deviation; * Ulceration in the mouth that healed without scarring, other mucosal areas were spared³²

490 **Table 2**

Table 2. Skin findings and configurations reported in cases of nonbullous cutaneous pemphigoid

Skin findings reported	No. of cases (%)
Erythematous, urticarial papules and plaques	69 (52.3%)
Papules/nodules	27 (20.5%)
Eczematous lesions	16 (12.1%)
No primary lesions reported¶	6 (4.5%)
Dermatitis herpetiformis-like lesions	5 (3.8%)
Ulcerations	3 (2.3%)
Erythroderma	3 (2.3%)
Other:	
Scarring alopecia	1 (0.8%)
Vegetations	1 (0.8%)
Solitary macule	1 (0.8%)
Excoriations	30 (22.7%)
Configuration reported	
Annular configuration*	8 (6.1%)
Figurated configuration	2 (1.5%)
Gyrated configuration	1 (0.8%)

491 The clinical presentation was reported in all 132 cases.

492 ¶ all 6 cases presented with secondary lesions in the form of excoriations

493 * two cases presented with erythema multiformis-like lesions

494

495 **Table 3**

496 **Table 3.** Reported localization of skin lesions in nonbullous cutaneous pemphigoid

Localization reported	No. of cases (%)
Extremities	43 (67.2%)
Trunk	42 (65.6%)
Generalized	14 (21.9%)
Head and/or neck	7 (10.9%)
Scalp	6 (9.4%)
Hands and/or feet	5 (7.8%)

498 The localization of the lesions was reported in 64 cases

499

500

501

502 **Table 4****Table 4.** Reported laboratory findings

	No. of cases with positive test results (%)	Reported in no. cases
DIF microscopy, linear IgG and/or C3c depositions along the BMZ	123 (93.2%)	132
IIF* IgG	42 (76.4%)	55
IIF on salt split skin, IgG, epidermal binding	46 (90.2%)	51
Nc16a ELISA, IgG	15 (57.7%)	26
BP230 ELISA, IgG	10 (52.6%)	19
Immunoblot BP180, IgG	11 (32.4%)	34
Immunoblot BP230, IgG	20 (55.6%)	36

DIF, direct immunofluorescence; IgG, immunoglobulin G; BMZ, basement membrane zone; IIF, indirect immunofluorescence; Nc16a, non-collagen 16a; ELISA, enzyme linked immunosorbent assay.

* different substrates were used by different authors.

503

504

505

506

507

508 **SUPPLEMENTARY MATERIAL**509 **Supplement 1**510 *Keywords used in the systematic search (performed in EMBASE & MEDLINE)*

511 ('non*bullous' AND 'pemphigoid') OR 'non*bullous pemphigoid' OR non*bullous bullous
 512 pemphigoid' OR 'non*bullous BP' OR 'pruritic pemphigoid' OR 'pruritic non*bullous
 513 pemphigoid' OR 'pemphigoid nodularis' OR 'nodular pemphigoid' OR 'prurigo nodularis-like
 514 pemphigoid' OR 'papular pemphigoid' OR 'prodromal BP' OR 'prodromal bullous
 515 pemphigoid' OR 'prodromal pemphigoid' OR 'prodrome of bullous pemphigoid' OR 'non
 516 bullous variant' NEAR/10 'pemphigoid' OR 'nonbullous variant' NEAR/10 'pemphigoid' OR
 517 'bullous pemphigoid mimicking' OR '-like bullous pemphigoid' OR 'erythrodermic bullous
 518 pemphigoid' OR ('bullous pemphigoid' AND 'without blister*') OR ('bullous pemphigoid'/exp
 519 AND 'without blister*') OR ('bullous pemphigoid' AND 'without bullae') OR ('bullous
 520 pemphigoid' AND 'without bullous lesions')


521

522

523 **Supplement 2**524 *List of all included articles presenting cases of nonbullous cutaneous pemphigoid*

First author	Publication year
Barker et al. ²⁰	1983
Bingham et al. ⁵⁸	1984
Amato et al. ²⁴	1988
Borradori et al. ⁵⁹	1990
Wolf et al. ⁶⁰	1992
Ross et al. ⁴⁹	1992
Strohal et al. ¹⁰	1993
Bourke et al. ⁴⁵	1994
Wever et al. ⁶¹	1995
Jeong et al. ⁶²	1995
Cliff et al. ⁶³	1996
Kawahara et al. ⁵³	1997
Alonso-Llamazares et al. ⁶⁴	1998
Alonso-Llamazares et al. ⁶⁵	1998
Scrivener et al. ⁶⁶	1999
Ameen et al. ³²	2000
Schmidt et al. ⁶⁷	2002
Powell et al. ⁶⁸	2002
Goel et al. ⁶⁹	2003
Mechtel et al. ⁷⁰	2003

Tashiro et al. ³⁰	2005
von Felbert et al. ⁷¹	2005
Lamb et al. ²¹	2006
Yesudian et al. ⁷²	2009
Matsudate et al. ⁷³	2009
Axelrod et al. ²⁵	2010
Safa et al. ⁷⁴	2010
McCourt et al. ⁷⁵	2010
Geiss Steiner et al. ⁷⁶	2010
Lehman et al. ⁷⁷	2011
Patel et al. ²⁹	2012
Bakker et al. ¹¹	2013
Balakirski et al. ⁷⁸	2014
Liu et al. ³¹	2014
Kabuto et al. ²⁷	2015
Altman et al. ²³	2015
Park et al. ²⁸	2015
Huet et al. ⁷⁹	2016
Ise et al. ²⁶	2016



Capsule summary

- What is already known on this topic.
Cutaneous pemphigoid can present without typical bullae and consequently diagnosis can be delayed.
- What this article adds to our knowledge.
The most frequently reported clinical features in these patients are pruritic, erythematous, urticarial papules and plaques.
- How this information impacts clinical practice and/or changes patient care.
Clinicians should consider nonbullous cutaneous pemphigoid in older patients with refractory pruritus.